

Identification of Capsular Serotypes K1 and K2 in Clinical Isolates of *Klebsiella Pneumoniae* in North of Iran

Rhokhsareh Akbari (MSc)

MSc student of Microbiology,
Department of Biology, Islamic Azad
University, Rasht Branch, Rasht, Iran

Leila Asadpour (PhD)

Assistant professor in Microbiology,
Department of Biology, Islamic Azad
University, Rasht Branch, Rasht, Iran

Corresponding author: Leila
Asadpour

Tel: +989113383860

Email: asadpour@iaurasht.ac.ir

Address: Islamic Azad University,
Rasht Branch, Rasht, Iran

Received : 03 Nov 2015

Revised: 01 Dec 2015

Accepted: 05 Dec 2015

ABSTRACT

Background and Objectives: *Klebsiella pneumoniae* is one of the most important nosocomial pathogens. Its capsular polysaccharide is considered as the first and most important virulence factor of this bacterium. This study aimed to investigate the presence of capsular serotypes K1 and K2 in *K. pneumoniae* isolates to examine the virulence potency of the isolates.

Methods: Overall, 65 capsulated *K. pneumoniae* isolates were collected from patients with urinary tract infections in Rasht, Iran. The isolates were examined using biochemical tests and CPS gene amplification using PCR. Mucoïd phenotype of the isolates was determined by the string test. The presence of K1 and K2 genes was evaluated by PCR using specific primers for the genes.

Results: Of 65 *K. pneumoniae* isolates, seven (10.77%) were positive for the presence of the K1 gene and four (6.15%) were positive for the presence of the K2 gene. In addition, six serotype K1 isolates (27.27%), four serotype K2 isolates (18.18%), and 12 non-K1/K2 serotype isolates (54.54%) had hypermucoviscosity phenotypes.

Conclusion: Our results confirm the presence of the capsular serotypes in *K. pneumoniae* isolates, with a relatively high prevalence for the capsular serotype K1. This study clarifies the importance of rapid diagnosis and suitable treatment of infections caused by *K. pneumoniae* in prevention of complicated infections.

Keywords: *Klebsiella pneumoniae*, Virulence factors, Capsular polysaccharide.

INTRODUCTION

Klebsiella pneumoniae is an opportunistic and important nosocomial pathogen that causes a wide range of infections including septicaemia, pneumonia, urinary tract infection, meningitis and abscesses in different organs, particularly liver (1). *Klebsiella* sp. are surrounded by a thick layer of hydrophilic capsular polysaccharide (CPS), which is considered as the first and most important virulence factor of these bacteria. CPS has an essential role in the attachment of bacteria to epithelial and mucosal surfaces. It also protects the bacteria from serum opsonization and phagocytosis, thereby hiding the bacteria from the host immune system. On the other hand, the capsule provides protection against adverse environmental conditions and reduces the permeability of antibiotics into the bacteria (2, 3). Capsular serotypes K1 and K2 strains of *K. pneumoniae* have more pathogenicity. These serotypes have shown more resistance against the bactericidal effect of serum and phagocytic ingestion compared to other serotypes (4). Therefore, the presence of K₁

and K₂ capsular serotypes of *K. pneumoniae* in clinical samples may be a sign of serious infections. The present study aimed at investigating the presence of K₁ and K₂ capsular serotypes of *K. pneumoniae* to examine the virulence potency of the clinical isolates of this bacterium.

MATERIAL AND METHODS

Overall, 65 *K. pneumoniae* isolates from patients with urinary tract infections were collected from clinical laboratories in city of Rasht, Iran. Bacterial genomic DNA was extracted using commercial DNA isolation kits for gram-negative bacteria (Cinnagen, Iran). All isolates were tested for the presence of *K. pneumoniae* capsular antigens by PCR, using CPS-specific primers (5). In addition, primers specific for capsular serotype K1- and K2-specifying genes were used in the PCR experiment. All PCR experiments were performed according to the method described by Feizabadi et al. (6). Table 1 shows the primers used in the PCR experiments.

Table 1: Oligonucleotide primers used for amplification of sequences specific for *K. pneumoniae* CPS, and K1 and K2 genes.


Gene	Primers	Product size (bp)
CPS	F: TATTCATCAGAAGCACGAGCTGGGAGAAGCC-3' R: GTCGGTAGCTGTAAAGCCAGGGGCGGTAGCG-3	418
K1	F: 5' ACGATAGAGGTGTATTGTCGC 3' R: 5' ACGATAGAG GTGTATTGTCGC3'	352
K2	R: 5' TGATACTTGACAGAGGGAGTA-5' F: 5' ACGATCGTTACAGTGACAAG-3'	321

RESULTS

In this study, all 65 *K. pneumoniae* isolates were gram-negative, non-motile, rod-shaped, lactose-fermenting, methyl-red negative, Voges-Proskauer positive, indol negative, catalase positive, urease positive and hydrogen sulphide negative, with mucoid colonies on MacConkey agar. The presence of *K. pneumoniae* CPS was confirmed by detecting


a 418 bp fragment after the PCR experiment (Figure 1). The specific primers yielded a 352 bp fragment in seven isolates (10.77%), which were identified as capsular serotype K1. In addition, a 321 bp fragment was produced in four isolates (6.15%), which were identified as capsular serotype K2. Moreover, 54 isolates (83.07%) were identified as non-K1/K2 serotypes (Figure 2).

Figure 1- Agarose gel electrophoresis of CPS gene PCR amplicons. Column M: 100bp DNA marker, Column 1: Positive control (418 bp amplicon of CPS confirmed by sequencing), Columns 2-6: CPS positive strains.



Medical Laboratory Journal, Jan, Feb 2017; Vol 11: No 1

Figure 2- Agarose gel electrophoresis of PCR amplicons for K1- and K2-specifying genes. Column M: 100bp DNA marker, Column 1: Positive control (352 bp amplicon of K1 confirmed by sequencing), Column 2: K1 positive strain, Column 3: Positive control (321 bp amplicon of K2 confirmed by sequencing), Column 4: K2 positive strain.



DISCUSSION

In the present study, 65 *K. pneumoniae* clinical isolates were examined to identify the K1 and K2 capsular serotypes. The results showed that seven isolates (10.77%) were serotype K1, while four (6.15%) were serotype K2. In a similar study in Iran, Feizabadi et al. (2013) reported the frequency of capsular serotypes K1 and K2 among 89 *K. pneumoniae* isolates from urine samples as 11.2% and 14.6%, respectively (6). Similar studies have been conducted in other parts of the world. In study of Yeh et al. on 50 *K. pneumoniae* clinical isolates from patients with liver abscesses in Taiwan, 26 isolates (52%) were identified as serotype K1 and 10 isolates (20%) were identified as serotype K2 (7). The results of these studies also showed that non-K1/K2 serotypes are involved in

aggressive *Klebsiella* infections. Another study in Taiwan by Liao et al. (2011) demonstrated that K1 is the most abundant serotype among *Klebsiella* isolates from cases of bacteraemia (8). In Iraq, study of Abdol-Razzaq et al. (2014) on 158 *Klebsiella* isolates found K1 and K2 capsular antigens in 18.6% and 32.5% of isolates, respectively. However, they found three cases that were positive for both serotypes (5). In a study by Tamara et al. on 40 *Klebsiella* isolates, 57.5% of isolates were identified as serotype K1, 27.5% were identified as serotype K2, and 15% were identified as non-K1/K2 serotypes (9). Study of Jiang Wang et al. on 101 *Klebsiella* isolates reported the prevalence of capsular serotypes K1, K2 and non-K1/K2 as 42.57%, 36.63% and 20.79%, respectively (10).

CONCLUSION

Capsular serotype K1 or K2 plays important roles in the pathogenesis of *K. pneumoniae* infections. Our study confirms the presence of these capsular serotypes in isolates from urine samples, with a relatively high prevalence for the serotype K1. This study clarifies the importance of rapid diagnosis and suitable treatment of infections caused by *K. pneumoniae* in prevention of complicated

REFERENCES

1. Yu W-L, Ko W-C, Cheng K-C, Lee H-C, Ke D-S, Lee C-C, et al. Association between *rmpA* and *magA* genes and clinical syndromes caused by *Klebsiella pneumoniae* in Taiwan. *Clinical Infectious Diseases*. 2006; 42(10): 1351-8.
2. Burmelle M, Bahl MI, Jensen LB, Sørensen SrJ, Hansen LH. Type 3 fimbriae, encoded by the conjugative plasmid *pOLA52*, enhance biofilm formation and transfer frequencies in *Enterobacteriaceae* strains. *Microbiology*. 2008; 154(1): 187-95.
3. Struve C, Bojer M, Nielsen EMI, Hansen DSd, Krogfelt KA. Investigation of the putative virulence gene *magA* in a worldwide collection of 495 *Klebsiella* isolates: *magA* is restricted to the gene cluster of *Klebsiella pneumoniae* capsule serotype K1. *Journal of medical microbiology*. 2005; 54(11): 1111-3.
4. Chhibber S, Aggarwal S, Yadav V. Contribution of capsular and lipopolysaccharide antigens to the pathogenesis of *Klebsiella pneumoniae* respiratory tract infection. *Folia microbiologica*. 2003; 48(5): 699-702.
5. Abdul-Razzaq MS, Al-Khafaji JKT, Al-Maamory EHK-a. Molecular characterization of capsular polysaccharide genes of *Klebsiella pneumoniae* in Iraq. *Int J Curr Microbiol App Sci*. 2014; 3(7): 224-34.
6. Feizabadi MM, Raji N, Delfani S. Identification of *Klebsiella pneumoniae* K1 and K2 capsular types by PCR and quellung test. *Jundishapur Journal of Microbiology*. 2013; 6(9): E7585. DOI: 10.5812/ijm.7585.
7. Yeh K-M, Kurup A, Siu L, Koh Y, Fung C-P, Lin J-C, et al. Capsular serotype K1 or K2, rather than *magA* and *rmpA*, is a major virulence determinant for *Klebsiella pneumoniae* liver abscess in Singapore and Taiwan. *Journal of clinical microbiology*. 2007; 45(2): 466-71.
8. Liao C-H, Huang Y-T, Lai C-C, Chang C-Y, Chu F-Y, Hsu M-S, et al. *Klebsiella pneumoniae* bacteremia and capsular serotypes, Taiwan. *Emerging infectious diseases*. 2011; 17(6): 1113. doi: 10.3201/eid1706.100811.
9. Zedan TH, Al-Jailawi MH, Jassim KA. Determination Of K1 And K2 Capsular Serotypes For *Klebsiella Pneumoniae* Using *Maga* And *K2a* Genes As Specific Molecular Diagnosis Tools.
10. Wang J, Yan Y, Xue X, Wang K, Shen D. Comparison of pyogenic liver abscesses caused by hypermucoviscous *Klebsiella pneumoniae* and non-*Klebsiella pneumoniae* pathogens in Beijing: a retrospective analysis. *J Int Med Res*. 2013 Aug;41(4):1088-97. doi: 10.1177/0300060513487645.

infections.

ACKNOWLEDGMENTS

The article is derived from an MSc thesis, financially supported by the Islamic Azad University of Rasht, Iran.

CONFLICT OF INTEREST

The authors declare no conflicts of interest.