

**Prevalence of cytogenetic and electrophoretic abnormalities in patients  
suspected of multiple myeloma**

**Running title:** Multiple myeloma and abnormalities on FISH

**Amudha Subramaniam**

St john's Medical College and Hospital Bangalore-560034, India- St john's  
Medical College and Hospital

Email: [amudha.s@stjohns.in](mailto:amudha.s@stjohns.in) ORCID: 0009-0004-5128-256X

**Saranya Raj**

Department of Biochemistry St john's Medical College and Hospital Bangalore-  
560034, India- St john's Medical College and Hospital

Email: [dr.saranyaraj.is@gmail.com](mailto:dr.saranyaraj.is@gmail.com)

**Thashreefa Olakara**

Department of Biochemistry St john's Medical College and Hospital Bangalore-  
560034, India- St john's Medical College and Hospital

Email: [thashreefa@gmail.com](mailto:thashreefa@gmail.com)

**Jayakumari S**

Department Biochemistry, St john's Medical College Sarjapur Road, Bengaluru  
560034

Email: [jayakumari.s@stjohns.in](mailto:jayakumari.s@stjohns.in) ORCID: 0000-0002-2104-8228

**Veronica Preetha Tilak**

Consultant Division of Molecular Biology and Genetics, St john's Medical  
College and Hospital Bangalore-560034, India- St john's Medical College and  
Hospital

Email: [Preetha444@yahoo.co.in](mailto:Preetha444@yahoo.co.in)

**Corresponding author:** Jayakumari S

**Email:** [jayakumari.s@stjohns.in](mailto:jayakumari.s@stjohns.in)

**Tel:** +919880259248

**Address:** Dr Jayakumari S, Department Biochemistry, St john's Medical  
College Sarjapur Road, Bengaluru 560034 Karnataka India.

## Abstract

**Background:** Multiple myeloma (MM) is a blood cancer characterized by the uncontrolled growth of plasma cells in the bone marrow. It typically develops from monoclonal gammopathy of undetermined significance (MGUS), which can progress to smoldering myeloma and eventually to symptomatic disease. Diagnosis is primarily established using serum protein electrophoresis (SPEP), immunofixation electrophoresis (IFE), and free light chain (FLC) testing. Additionally, Fluorescence in Situ Hybridization (FISH) plays a crucial role in identifying genetic abnormalities that influence the disease course and prognosis. This study aims to evaluate the prevalence of electrophoretic and genetic abnormalities among patients referred for serum protein electrophoresis, with a focus on cytogenetic abnormalities detected by FISH in confirmed MM cases.

**Methods:** Samples received for SPEP from 2017 to 2023 were analysed. Patients with abnormalities on electrophoresis (such as distortions or M-spikes) underwent further evaluation, including immunofixation, free light chain assays, bone marrow examination, and other hematologic investigations. Confirmed MM cases were referred for FISH analysis to identify common cytogenetic abnormalities

**Results:** Out of 800 patients with electrophoretic abnormalities, 100 were confirmed to have multiple myeloma. FISH analysis was available for 68 of these cases, detecting cytogenetic abnormalities in 67.6% of patients. The most common abnormalities were IGH break apart (54.5%), followed by p53 deletion (23.5%), t (4; 14) (14.7%), t (14q24;q20) (7.4%), monosomy 13 (5.9%), and monosomy 14 (4.4%).

**Conclusion:** The majority of MM patients showed abnormalities on FISH, with IGH break-apart being the most frequently detected. These cytogenetic abnormalities provide valuable prognostic information and can help guide treatment decisions. This study emphasizes the importance of routine cytogenetic profiling in MM to optimize therapeutic outcomes. Molecular cytogenetic techniques, especially FISH, are essential tools in evaluating suspected multiple myeloma. They play a pivotal role in the detection of genetic abnormalities, guiding treatment strategies, and ultimately improving patient outcomes.

**Keywords:** Multiple myeloma, Electrophoresis, Fluorescence in Situ Hybridization (FISH)

## Introduction

Multiple myeloma (MM) is a cancer involving plasma cells—specialized white blood cells that produce antibodies. The disease is defined by excessive, uncontrolled growth of a single clone of plasma cells within the bone marrow. This abnormal expansion results in increased production of monoclonal immunoglobulins or light chains, which can ultimately lead to damage in various organs. MM develops from a precursor condition known as monoclonal gammopathy of undetermined significance (MGUS), which may progress to smoldering myeloma and eventually to symptomatic MM (1). It often presents with bone pain, anaemia, hypercalcemia, and renal dysfunction, but diagnosis requires specific laboratory evaluations (2). SPEP, IFE, and FLC analysis are the standard methods for MM diagnosis (3). In addition, the identification of genetic abnormalities using FISH has provided valuable information about the disease course and treatment response (4). Several key abnormalities are routinely studied due to their prognostic implications. These translocations, present in 10-20% of patients, influence prognosis, with t(4;14) associated with poor outcomes, while t(11;14) has a more neutral risk profile (5). Deletion of 17p (TP53): Loss of the TP53 gene located on chromosome 17p is a major high-risk abnormality, strongly linked to more aggressive disease behavior and poorer patient survival. This abnormality occurs in 10-20% of cases and is a critical indicator of poor prognosis (6). Monosomy 13 or 13q Deletion: Deletion of chromosome 13 or loss of genetic material at 13q14 is seen in about 50% of MM patients. Although historically considered a poor prognostic factor, its impact on outcomes has been overshadowed by other markers (7). Gain/amplification of CKS1B on 1q21: Amplification of the 1q21 is seen in up to 40% of MM cases and is associated with worse outcomes, increased relapse rates, and shorter survival (8). t(14;20) and t(14;16): these rare translocations are observed in 1-2% and 5-10% of MM patients, respectively, and are considered high-risk features that lead to poorer prognosis and shorter overall survival (9). These acquired cytogenetic abnormalities in MM samples are considered as “known biomarkers” which assist in evaluating prognosis and determining how well a patient may respond to therapy. The importance in forecasting prognosis and informing therapy, these cytogenetic abnormalities are being considered into risk stratification guidelines (10). Cytogenetic methods are essential for detecting chromosomal abnormalities, studying genome structure, and identifying genetic disorders and malignancies (11). FISH is a sensitive molecular cytogenetic technique used to detect chromosomal abnormalities with high resolution, even at the single-gene level (12). Specific recommendations for performing FISH in myeloma by The European Myeloma Network aims to build a proper understanding of the pathogenesis of Myeloma and to provide standard approaches using specific Probes to establish a common cytogenetic biomarker for diagnostic work-up and there by classifying the disease into subgroups with prognostic and predictive outcomes (13).

Despite significant recent developments in the detection and therapy of MM, the disease remains highly heterogeneous, with variable clinical outcomes based on genetic and molecular abnormalities. Understanding the genetic profile of MM is critical for tailoring treatment strategies and improving patient outcomes. While extensive research has identified several key cytogenetic abnormalities associated with MM prognosis, the prevalence and distribution of these abnormalities can vary across populations and geographic regions and data regards same is limited in literature. This study aims to evaluate the prevalence of electrophoretic and cytogenetic abnormalities in patients referred for serum protein electrophoresis in our population. Understanding these genetic abnormalities enables personalized treatment strategies, with high-risk patients benefiting from more aggressive therapeutic approaches.

## Methods

This study is retrospective study. All serum electrophoresis samples received between January 2017 and December 2023 in the Department of Biochemistry at St John's Medical College were included. A total of 800 patients showing distortions, M-band, or spikes on SPEP were further evaluated using additional tests, including free light chain assays, immunofixation, and bone marrow studies.

Cytogenetic analysis was carried out at Division of Human Genetics, St. John's Medical College, for patients confirmed with MM. FISH was performed on whole marrow samples to detect cytogenetic abnormalities specific to MM, using a comprehensive FISH panel of probes from Metasystems, Germany. These probes targeted various genes and chromosomal regions implicated in MM, such as the IGH break apart probe on 14q32.3, and TP53 on 17p13.

FISH probes were applied on fixed cell suspension following standardized FISH protocol which included probe addition, co-denaturation at 75°C for 2 minutes, and hybridization at 37°C using Euroclone. Post hybridization washes at 72°C, followed by dehydration in ethanol series and DAPI addition. 200 nuclei are analysed using BX53F Olympus fluorescent microscope and digital imaging.

### **FISH Probe Specifications from XCyting FISH Probes of MetaSystems**

1. IGH BA:- Orange-labelled probe partly binds to constant region of the IGH locus at 14q32.3 and a green-labelled probe hybridizing to the variable distal region of the IGH locus at 14q32.3 (Fig 1).
2. t (11;14) MYEOV/IGH DF:- Orange-labelled probe binds to the MYEOV/CCND1 gene region at 11q13.3 and a green-labelled probe hybridizing to the IGH gene region at 14q32.3 (Fig 2).
3. t (4;14) FGFR3/IGH DF:- Orange-labelled probe binds to the FGFR3 gene region at 4p16.3 and a green-labelled probe hybridizing to the IGH gene region at 14q32.3 (Fig 3).
4. t (14;16) IGH/MAF DF:- Green-labelled probe binds to the IGH gene region at 14q32.3 and an orange-labelled probe hybridizing to the WWOX/MAF gene region at 16q23 (Fig 4).
5. DLEU/LAMP:- Orange-labelled probe binds to the DLEU1/MIR15A/MIR16-1 gene region at 13q14.2 including D13S319 and a green-labelled probe hybridizing to the LAMP1 gene region at 13q34.
6. TP53/NF1 :-Orange-labelled probe binds to the TP53 gene region at 17p13 and a green-labelled probe hybridizing to the NF1 gene region at 17q11.2.
7. t (6;14) CCND3/IGH DF :- Orange-labelled probe binds to the CCND3 gene region at 6p21.1 and a green-labelled probe hybridizing to the IGH gene region at 14q32.3
8. t (14;20) IGH/MAFB DF :- Green-labelled probe binds to the IGH gene region at 14q32.3 and an orange-labelled probe hybridizing proximal to the MAFB gene region 20q12.

Data from confirmed MM cases and their FISH results were collected and analysed. Frequencies of cytogenetic abnormalities were calculated, and correlations between genetic findings and disease characteristics were assessed by chi-square tests.

## Results

Eight hundred patients were evaluated for electrophoretic abnormalities. Out of these, 100 patients were diagnosed with MM, and 68 of these underwent FISH testing (Table 1).

**Genetic Abnormalities Detected:** 200 interphase cells were analysed for each of the probe applied. Among the 68 MM patients with FISH data, 67.6% exhibited cytogenetic abnormalities. The most common abnormality was the IGH break-apart (54.5%), followed by TP53 deletion. Further FISH analysis on positive cases with IGH break-apart, t(4;14), and

t(14;20) translocations were observed to be more evident. Additional findings included monosomy 13/deletion 13q and monosomy 14/deletion 14q32.2 (table 2).

## Discussion

This study demonstrates that a significant proportion (67.6%) of MM patients exhibit cytogenetic abnormalities detectable by FISH. The most prevalent abnormality, IGH break-apart, was found in 54.5% of cases. This abnormality is well-known for its association with MM and is linked to has been linked to unfavorable outcomes in multiple studies. The detection of TP53 deletion in 23.5% of cases also aligns with existing literature, as TP53 abnormalities often correlate with aggressive disease and reduced survival.

Translocations such as t(4;14) and t(14;20), while less common, were also detected in our cohort. These translocations have been implicated in high-risk disease and poorer response to conventional therapies. Monosomy 13 and monosomy 14, although infrequent, may contribute to disease progression and should be evaluated in larger studies. Several previous studies have reported similar patterns of genetic abnormalities in MM, although variations in the prevalence of specific abnormalities can be observed depending on the population and methodologies used. Our finding that IGH break-apart was present in 54.5% of cases is consistent with other studies, which typically report IGH rearrangements in 40-60% of MM patients. Rajkumar et al, found a prevalence of 50% for IGH abnormalities, which are known to be associated with a poor prognosis and resistance to conventional therapy (2). Similarly, Fonseca et al, found that IGH translocations were the most common cytogenetic abnormality in MM, occurring in around 50% of cases, emphasizing its role in MM pathogenesis. TP53 Deletion (23.5%): The prevalence of p53 deletion in our study (23.5%) is slightly higher compared to some studies, where the frequency typically ranges between 10-20% (14).

According to Amudha et al., the translocations t(4;14), t(14;16), t(6;14), and t(14;20) showed an association with anemia. The t(4;14) abnormality, in particular, was linked to elevated serum monoclonal protein levels and increased plasma cell proliferation. In addition, monosomy 13 has been correlated with reduced survival and a tendency for progression from monoclonal gammopathy (15). Other studies such as Abdallah et al have reported that TP53 deletions were detected in approximately 17% of MM patients and were strongly associated with advanced disease and poorer prognosis. This higher percentage in our cohort may reflect a referral bias of advanced or aggressive MM cases. t(4;14) Translocation (14.7%): The translocation t(4;14), detected in 14.7% of patients, is a well-established high-risk feature in MM. Reported frequencies for this abnormality in other studies range from 10-15%. similarly, t(4;14) were reported in 15% of their cohort and noted that this translocation was associated with poor prognosis and a lower response to standard treatments (16). In contrast, Fonseca et al, reported a prevalence of around 13%, which further supports our findings. t (14;20) Translocation (7.4%): Although t(14;20) is considered a rare genetic abnormality in MM, our study found a prevalence of 7.4%. This is in line with reported frequencies of 5-7% in previous studies. Patients with this translocation are often considered to have high-risk disease with a shorter overall survival Fonseca et al (14). Monosomy 13 and 14: The detection of monosomy 13 (5.9%) and monosomy 14 (4.4%) in our study is consistent with other findings. Rajkumar et al, found that monosomy 13 occurs in about 15% of newly diagnosed MM patients and is associated with aggressive disease (2). However, our slightly lower detection rate may reflect differences in the specific populations being studied, or the fact that other genetic factors are now recognized as more predictive of outcomes in MM.

Amare et al. reported that FISH is a valuable and straightforward technique for detecting a wide range of cytogenetic abnormalities, providing important insights for disease risk stratification. They also noted that, when combined in the future with mutational and gene-

expression profiling, FISH could contribute to more precise disease characterization and help identify actionable therapeutic targets (17).

This study highlights the importance of electrophoretic and genetic testing in the diagnosis and prognosis of MM. The prevalence of genetic abnormalities in MM patients, especially IGH break-apart, provides critical information about disease characteristics and expected outcomes. Further research is required to explore the therapeutic implications of these findings, particularly in the context of newer targeted therapies.

## **Conclusion**

This study, along with previous research, emphasizes the critical importance of comprehensive cytogenetic profiling in MM patients. FISH analysis remains a valuable tool for identifying key genetic abnormalities that can influence prognosis and guide treatment decisions. Further studies exploring these abnormalities in different populations and their response to emerging therapies are necessary to refine prognostic models and optimize MM management strategies. The prevalence of IGH break-apart and TP53 deletions underscores their importance as biomarkers for disease aggressiveness and resistance to therapy. Translocations such as t(4;14) and t(14;20), though less common, highlight high-risk disease profiles that may benefit from early intervention with novel therapies. The molecular cytogenetic techniques, especially FISH, are indispensable tools in the evaluation of suspected multiple myeloma. They play a pivotal role in the detection of genetic abnormalities, guiding treatment strategies, and ultimately improving patient outcomes.

A limitation of this study is the relatively small sample size. The samples were analysed using probes that are limited to specific variations, and therefore, variations not covered by these probes cannot be detected. The ideal method for such cases would have been the use of mFISH and SKY. A future multicentric study on cytogenetics evaluation with electrophoresis on a larger population case with MGUS, SMM and Multiple MM is recommended.

## **Acknowledgment**

We extend our gratitude to everyone whose contributions have been invaluable to this study.

## **Funding sources**

Not applicable.

## **Ethical statement**

Institutional Ethics Committee approval for the study was obtained (IEC reference number 297/2022). A waiver of consent was obtained. There is no direct contact between the researcher and participant.

## **Conflict of interest**

The authors declare that they have no competing interests.

## **Author contributions**

Amudha Subramaniam Interpreted FISH laboratory reports. Saranya Raj contributed to the selection of lab and clinical data. Thashreefa Olakara reviewed medical records, collected and monitored data, interpreted results, maintained patient files and the master file, drafted the final report. Jayakumari S contributed to the concept and design, interpreted laboratory reports (electrophoresis), monitored data, performed statistical analysis and interpretation, drafted the final report. Veronica Preetha Tilak Interpreted FISH laboratory reports.

## Data availability statement

The data that support the findings of this study are available from the corresponding author upon reasonable request.

## References

1. Palumbo A, Anderson K. Multiple myeloma. *N Engl J Med*. 2011;364(11):1046-60.
2. Rajkumar SV, Kumar S. Multiple myeloma: diagnosis and treatment. *Mayo Clin Proc*. 2016;91(1):101-19.
3. Willrich MA, Katzmann JA. Laboratory testing requirements for diagnosis and follow-up of multiple myeloma and related plasma cell dyscrasias. *Clin Chem Lab Med*. 2016;54(6):907-19.
4. Ratan ZA, Zaman SB, Mehta V, Haidere MF, Runa NJ, Akter N. Application of fluorescence in situ hybridization (FISH) technique for detection of genetic aberration. *Cureus*. 2017;9(6):e1325.
5. Puertas B, González-Calle V, Sobejano-Fuertes E, Escalante F, Rey-Bua B, Padilla I, et al. Multiple myeloma with t(11;14): impact of novel agents on outcome. *Blood Cancer J*. 2023;13(1):40.
6. Campo E, Cymbalista F, Ghia P, Jäger U, Pospisilova S, Rosenquist R, et al. TP53 aberrations in chronic lymphocytic leukemia. *Haematologica*. 2018;103(12):1956-68.
7. Fonseca R, Oken MM, Harrington D, Bailey RJ, Van Wier SA, Henderson KJ, et al. Deletions of chromosome 13 in multiple myeloma. *Leukemia*. 2001;15(6):981-6.
8. Hanamura I. Gain/amplification of chromosome arm 1q21 in multiple myeloma. *Cancers*. 2021;13(2):256.
9. Schavougoulidze A, Perrot A, Cazaubiel T, Leleu X, Montes L, Jacquet C, et al. Prognostic impact of t(14;16) in multiple myeloma. *Blood Cancer J*. 2023;13:160.
10. Lu X, Andersen EF, Banerjee R, Eno CC, Gonzales PR, Kumar S, et al. Guidelines for cytogenetic testing/reporting. *Blood Cancer J*. 2025;15(1):86.
11. Shakoori AR. Fluorescence in situ hybridization and its applications. In: *Chromosome Structure and Aberrations*. 2017. p. 343-67.
12. Chester M, Leitch AR, Soltis PS, Soltis DE. Application of modern cytogenetic methods. *Genes*. 2010;1(2):166-92.
13. Ross FM, Avet-Loiseau H, Ameye G, Gutiérrez NC, Liebisch P, O'Connor S, et al. EMN report on interphase FISH in multiple myeloma. *Haematologica*. 2012;97(8):1272-7.
14. Fonseca R, Bergsagel PL, Drach J, Shaughnessy J, Gutierrez N, Stewart AK, et al. IMWG molecular classification of multiple myeloma. *Leukemia*. 2009;23(12):2210-21.
15. Amudha S, Drishya R, Veronica PT, Raina M, Mary MA. Cytogenetics of multiple myeloma. *Indian J Genet Mol Res*. 2023;12.
16. Abdallah N, Rajkumar SV, Greipp P, et al. Cytogenetic abnormalities in multiple myeloma. *Blood Cancer J*. 2020;10:82.
17. Amare PK, Khasnis SN, Hande P, Lele H, Wable N, Kaskar S, et al. Cytogenetic abnormalities in multiple myeloma. *Cytogenet Genome Res*. 2023;162(10):529-40.


**Table 1.** Distribution of electrophoretic and genetic findings

Parameter	Number of patients	Percentage (%)
Total patients evaluated by SPEP	800	100
Confirmed cases of MM	100	12.5
Patients with FISH data available	68	68
Patients with cytogenetic abnormalities	46	67.6




**Table 2.** Prevalence of genetic abnormalities in MM


<b>..Genetic Abnormality</b>	<b>Frequency (%)</b>	<b>International Myeloma Society/International Myeloma Working Group (IMWG) (10).</b>	<b>Risk stratification</b>
IGH break-apart	54.5	40% of MM can be defined IGH- break-apart IGH rearrangement are always associated with either t(4;14) or t(6;14) or t(14;20) or t(11;14) or t(14;16)	High Risk
TP53 deletion	23.5	80%	High Risk
t(4;14) translocation	14.7	40% of IGH-Break-apart are followed with t(4;14) (p16; q32)	High Risk
t(14;20) translocation	7.4	40% of IGH-Break-apart are followed with t(14;20)(q32; q12),	High Risk
Monosomy/Deletion 13q	5.9	All most all cases will have 13q deletion	Standard Risk
Monosomy/deletion 14q32.2	4.4	Biallelic deletion of IGH gene on 14q32.2	High Risk




**Figure 1.** Nuclei showing IGH break-apart on 14q32.3 Probe Specifications: IGH BA consists of an orange-labelled probe partly covering the constant region of the IGH locus at 14q32.3 and a green-labelled probe hybridizing to the variable distal region of the IGH locus at 14q32.3.



**Figure 2.** t(11;14) MYEOV/IGH DF is designed as a dual fusion probe. The orange labelled probe hybridizes to region 11q13 including CCND1, the green labelled probe flanks the IGH breakpoint region at 14q32



**Figure 3.** Probe specification. t(4;14) FGFR3/IGH DF consists of an orange-labelled probe hybridizing to the FGFR3 gene region at 4p16.3 and a green-labelled probe hybridizing to the IGH gene region at 14q32.3. Nuclei showing extra fusion signals for t(4;14) translocation



**Figure 4.** t(14;16) IGH/MAF DF consists of a green-labelled probe hybridizing to the IGH gene region at 14q32.3 and an orange-labelled probe hybridizing to the WWOX/MAF gene region at 16q23. Nuclei showing fusion signals for t(14;16) translocation.